

Edited date: 2020-06-01

Kidney transplantation- Metabolic syndrome

1. Lee MC, Lee CJ, Ho GJ, Lee CC, Shih MH, Chou KC, Hsu BG*. Hyperleptinemia positively correlated with metabolic syndrome in renal transplant recipients. *Clin Transplant*. 2010; 24:E124-9. (SCI)
2. Lee MC, Ho GJ, Chen JL, Hsu BG*. C-reactive protein positively correlates with metabolic syndrome in kidney transplantation patients. *Tzu Chi Med J*. 2010; 22:131-6.
3. Lee MC, Lee CC, Ho GJ, Chou KC, Shih MH, Hsu BG*. Hypoadiponectinemia correlates with metabolic syndrome in kidney transplantation patients. *Transplant Proc*. 2011; 43:2601-5. (SCI)
4. Hsu BG, Shih MH, Yang YC, Ho GJ, Lee MC*. Fasting long-acting natriuretic peptide correlates inversely with metabolic syndrome in kidney transplantation patients. *Transplant Proc*. 2012; 44:646-50. (SCI)
5. Huang IC, Chang CC, Lee CJ, Chen YC, Lee MC*, Hsu BG*. Positive correlation of serum adipocyte fatty acid binding protein levels with metabolic syndrome in kidney transplantation patients. *International Journal of Clinical and Experimental Pathology* 2017;10(8):8727-8734. (SCI) (*co- corresponding author)
6. Lee KM[¶], Lee MC[¶], Lee CJ, Chen YC, Hsu BG*. Inverse association of N-terminal pro-B-type natriuretic peptide level with metabolic syndrome in kidney transplantation patients. *Transplant Proc*. 2018; 50(8):2496-2501. (SCI) (¶
co-first author)
7. Chen PC¹, Chang YD¹, Lee MC*, Hsu BG*. High serum fibroblast growth factor 23 level is associated with metabolic syndrome in kidney transplantation patients. *Transplant Proc*. 2020;S0041-1345(20)30351-1. (SCI) (*co-corresponding author)

Kidney transplantation- Bone mineral disease

1. Hsu BG, Ho GJ, Lee CJ, Yang YC, Chen YC, Shih MH, Lee MC*. Inverse association of serum long-acting natriuretic peptide and bone mineral density in renal transplant recipients. *Clin Transplant*. 2012; 26(2):E105-10. (SCI)
2. Lee MC, Lee CJ, Shih MH, Ho GJ, Chen YC, Hsu BG*. N-terminal pro-B-type natriuretic peptide is inversely related to bone mineral density in renal transplant recipients. *Transplant Proc*. 2014; 46(10):3443-7. (SCI)
3. Hsu BG, Chen YC, Ho GJ, Shih MH, Chou KC, Lin TY, Lee MC*. Inverse association between serum osteoprotegerin and bone mineral density in renal transplant recipients. *Transplant Proc*. 2016; 48(3):864-69. (SCI)

Kidney transplantation- Arterial stiffness

1. Chen YC[¶], Lee MC[¶], Lee CJ, Ho GJ, Yin WY, Chang YJ, Hsu BG*. N-terminal pro-B-type natriuretic peptide associated with arterial stiffness by cardio-ankle vascular index in renal transplant recipients. *J AtherosclerThromb.*2013;

20(7):646-53. (SCI) (

¶

co-first author)

2. Ho GJ, Chen YC, Yin WY, Chang YJ, Lee MC*, Hsu BG*. Fasting serum long-acting natriuretic peptide correlates with the ankle brachial index in renal transplant recipients.

(Co-correspondence)) *Exp Clin Transplant.* 2013; 11(4):303-9. (SCI) (*co- corresponding author)

3. Lee MC, Chen YC, Ho GJ, Shih MH, Chou KC, Hsu BG*. Serum leptin levels positively correlate with peripheral arterial stiffness in kidney transplantation patients. *Transplant Proc.*2014; 46(2):353-8. (SCI)

4. Ho GJ[¶], Lee MC[¶], Lee JC, Chen YC, Hsu BG*. Hypoadiponectinemia correlates with arterial stiffness in kidney transplantation patients. *Clin Exp Nephrol.*

2015; 19(3):534-41. (SCI) (

¶

co-first author)

5. Hsu BG, Shih MH, Chen YC, Ho GJ, Lin TY, Lee MC*. High serum osteoprotegerin is associated with arterial stiffness in kidney transplant patients. *Tohoku J. Exp. Med.* 2015; 236(4): 247-53. (SCI)

6. Lee MC, Lee CJ, Hsu BG*. Serum osteopontin level correlates with carotid-femoral pulse wave velocity in kidney transplantation patients. *Acta Nephrologica.* 2015; 29(2): 86-93.

7. Tsai JP[¶], Lee MC[¶], Chen YC, Ho GJ, Shih MH, Hsu BG*. Hyperleptinemia is a risk factor at the development of central arterial stiffness in kidney transplant patients.

Transplant Proc

. 2015; 47(6):1825-30. Erratum in:

Transplant Proc

. 2015 Oct;47(8):2556. Le, M C [Corrected to Lee, M C]. (SCI) (

¶

co-first author)

8. Hsu BG, Lee CJ, Chen YC, Ho GJ, Lin TY, Lee MC*. Serum osteoprotegerin levels associated with aortic augmentation index in renal transplant recipients. *Tzu Chi Med J.* 2016; 28:20-23.

9. Ho CC, Hsu BG, Yin WY, Ho GJ, Chen YC, Lee MC*. Serum adiponectin is a negative predictor of central arterial stiffness in kidney transplant patients. *Scand J Clin Lab Invest* 2016; 76(3):264-269. (SCI)

10. Hsu BG, Liou HH, Lee CJ, Chen YC, Ho GJ, Lee MC*. Serum Sclerostin as an Independent Marker of Peripheral Arterial Stiffness in Renal Transplantation Recipients-A Cross-Sectional Study. *Medicine (Baltimore).* 2016; 95(15):e3300. (SCI)

11. Hsu BG, Lee CJ, Chen YC, Ho GJ, Lin TY, Lee MC*. Low serum-free oxygen radicals defense level is associated with peripheral arterial stiffness in kidney transplantation patients. *In t J Clin Exp Pathol*

2016; 9(10):10698-10706.(SCI)

12. Chen YC[¶], Hsu BG[¶], Ho CC, Lee CJ, Lee MC*. Elevated serum osteoprotegerin may predict peripheral arterial disease after kidney transplantation: A single-center prospective cross-sectional study in Taiwan. *PeerJ* 2017; 5:e3847<https://doi.org/10.7717/peerj.3847> (SCI) (

¶
co-first author)

13. Chen YC, Hsu BG, Lee CJ, Ho CC, Ho GJ, Lee MC*. Serum adipocyte fatty acid binding protein levels is associated with peripheral arterial stiffness quantified by cardio-ankle vascular index in renal transplant patients. *Clin Exp Nephrol* 2018; 22:188-195. (SCI)

14. Chen YC[¶], Lee MC[¶], Lee CJ, Hsu BG*. Hyperleptinemia is associated with aortic augmentation index in kidney transplant recipients. *Tzu Chi Medical Journal* 2018; 30(3):152-7. (

¶
co-first author)

15. Chen TL[†], Lee MC[†], Ho CC, Hsu BG*, Tsai JP*. Serum adipocyte fatty acid binding protein level is negatively associated with vascular reactivity index measured by digital thermal monitoring in kidney transplant patients. *Metabolites* 2019, 9(8), 159; <https://doi.org/10.3390/metabo9080159>. (SCI) (†co-first author, *co-corresponding author)

16. Lai YH[†], Lee MC[†], Ho GJ, Liu CH*, Hsu BG*. Association of low serum L-carnitine levels with peripheral arterial stiffness in patients who undergone kidney transplantation. *Nutrients* 2019, 11(1), 159; doi:10.3390/nu11092000 (SCI) (†co-first author, *co-corresponding author)

17. Liu YJ¹, Tsai JP¹, Wang LH, Lee MC*, Hsu BG*. Positive correlation of serum fibroblast growth factor 23 with peripheral arterial stiffness in kidney transplantation patients [published online ahead of print, 2020 Feb 15]. *Clin Chim Acta*. 2020;S0009-8981(20)30069-3. doi:10.1016/j.cca.2020.02.014 (SCI) (*co-corresponding author)

Kidney transplantation- Posttransplant malignancy

1. Chen JL, Ho GJ, Yang YC, Shih MH, Chou KC, Lee MC*. High incidence of transitional cell carcinoma in kidney transplant recipients in eastern Taiwan. *Tzu Chi Med J*. 2009; 21:118-22.

2. Yin WY, Koo M, Lee MC, Lu MC*. Incidence of non-Hodgkin lymphomas and the ten most commonly diagnosed cancers after heart transplantation: A nationwide population-based study in Taiwan. *Transplantation*. 2014; 98(7):e71-e73. (SCI)

Kidney transplantation- BK virus

1. Lai NS, Lu MC, Lee MC, Lin TY, Yin WY*. Detection of polyomavirus JC genotype from transplant patients by capillary electrophoresis: Comparison to fragment length polymorphism analysis. *J Formos Med Assoc*. 2008; 107(3):239-44. (SCI)

2. Yin WY, Lu MC, Lee MC, Liu SC, Lin TY, Lai NS*. A correlation between polyomavirus JC virus quantification and genotypes in renal transplantation. *Am J Surgery*. 2010; 200:53-8. (SCI)

3. Lee MC, Lu MC, Lai NS, Liu SC, Yu HC, Lin TY, Hung SP, Huang HB, Yin WY*. Renal dysfunction by BK virus infection is correlated with activated T cell level in renal transplantation. *J Surg Res*

□□□□

□□□ □□□□□□

□□, 01 □□ 2020 09:17 - □□□□ □□, 19 □□ 2020 09:00
